

Dynamical Properties of Quantum Hall Edge States

A.V.Balatsky⁺* and S.I.Matveenko ^{*+}

⁺ *Theoretical Division, Los Alamos National Laboratory, Los Alamos and, New Mexico 87545*

^{*} *Landau Institute for Theoretical Physics, Moscow, Kosygina str., 2, Russia, 117940*

(April 5, 1995)

Abstract

We consider the dynamical properties of simple edge states in integer ($\nu = 1$) and fractional ($\nu = 1/2m + 1$) quantum Hall (QH) liquids. The influence of a time-dependent local perturbation on the ground state is investigated. It is shown that the orthogonality catastrophe occurs for the initial and final state overlap $|\langle i|f \rangle| \sim L^{-\frac{1}{2\nu}(\frac{\delta}{\pi})^2}$ with the phase shift δ . The transition probability for the x-ray problem is also found with the index, dependent on ν . Optical experiments that measure the x-ray response of the QH edge are discussed. We also consider electrons tunneling from one dimensional Fermi liquid into a QH fluid. For any filling fraction the tunneling from a Fermi liquid to the QH edge is suppressed at low temperatures and we find the nonlinear $I - V$ characteristics $I \sim V^{1/\nu}$.

The quantum Hall (QH) liquid is an incompressible state, where all bulk excitations have finite energy gaps. As for any incompressible liquid, droplet of water being the simplest example, the only low energy excitations are surface modes. Upon quantizing these surface modes one arrives at the picture of the simple propagating excitations on the surface of the droplet. For the two dimensional QH liquid these excitations represent the extended gapless

edge excitations [1,2]. As was shown by Halperin, edge excitations in the integer QH (IQH) systems are described by a chiral 1D Fermi liquid theory [1]. Similarly, Wen showed that in the fractional QH (FQH) state the edge excitations are described by a chiral Luttinger liquid theory [2]. Within last few years the transport in the chiral Luttinger liquid was investigated in great detail, see for example [2,3]. We argue that the time dependent phenomena in this system deserve a separate detailed investigation as well.

In this article few dynamical phenomena in the QH edge state will be addressed. The simplest question reflecting the nontrivial dynamics of QH state is the orthogonality catastrophe [4]. We calculate the response of the QH edge states to local time-dependent perturbations. And show that, for the case of sufficiently fast switching on of a perturbation potential or slow switching off, the orthogonality catastrophe takes place. We find that the filling factor ν enters into nontrivial exponents as well as in the phase shift $\delta = -\nu V_0/(2v)$, where V_0 is the local potential, and v is the velocity of the edge mode (in 1D the density of states $N_0 = 1/v$). In the case of the x-ray problem the transition rate is given by a power law of the frequency near the threshold $W(\omega) \sim (\omega + E_0)^\gamma$ with power $\gamma = (1 - \delta/\pi)^2/\nu - 1$, where E_0 is the deep core hole energy.

We also consider the tunneling via a point contact of the electron from the metal into the QH edge. The strongly correlated nature of the QH liquid with the composite boson order parameter, representing binding of the flux with the fermions makes the excitations in the QH states sufficiently different from the quasiparticles in a Fermi liquid. This results in the suppression of the tunneling conductance at low temperature, bias voltage or driving frequency. We find $I \sim V^{2/\nu+1}$ for the tunneling characteristics.

The orthogonality catastrophe and x-ray singularity were previously considered for another 1-D system: The Luttinger liquid [5]. There are qualitative differences between the Luttinger liquid and the QH edge response to the x-ray transition. Firstly, the backscattering, important for the x-ray response of a Luttinger liquid, is not present in the QH system due to chiral nature of the state (we do not consider the role of disorder and the transition between opposite edges). Secondly, the presence of the fractionally charged quasiparticles

in the QH liquid leads to the “fractionalization” of the phase shift with ν entering into all relevant exponents in the problem. Although natural for the QH state, this nontrivial dependence on the filling fraction (i.e. fractional statistics and fractional charge) can not be observed in the conventional Luttinger liquid.

We consider fractional states at odd integer ν^{-1} . In this case the edge state is known to be a single channel chiral Luttinger liquid [2] and the approach, used in [2,3], will be used.

Local time-dependent perturbation. The Hamiltonian of the edge waves is given by [2]

$$H = \int dx \frac{\pi}{\nu} \rho^2 v = \frac{2\pi}{\nu} v \sum_{k>0} \rho_k \rho_{-k} \quad (1)$$

where ρ is the one dimensional density, $v = cE/B$ is the velocity of electron drift near the edge, E is the electric field of the potential well, B is the magnetic field. The Hamiltonian Eq.(1) is quantized in terms of the $U(1)$ Kac-Moody algebra [2] and we have

$$[\rho_k, \rho_{k'}] = \frac{\nu}{2\pi} k \delta_{k+k',0} \quad [H, \rho_k] = v k \rho_k \quad (2)$$

A similar algebra (with change $\nu \rightarrow L$) is known in the Tomonaga model [6]. The electron operator is written in terms of the field φ as $\Psi \propto \exp i\varphi/\nu$ with the commutator

$$[\varphi(x), \varphi(y)] = i\pi\nu \text{sign}(y - x) \quad (3)$$

The density operator $\rho(x)$ is equal to $\rho(x) = \partial_x \varphi / (2\pi)$. The action in terms of the chiral boson field φ is

$$S = \frac{1}{4\pi\nu} \partial_x \varphi (\partial_t \varphi + v \partial_x \varphi) \quad (4)$$

The interaction of the system with an external time-dependent perturbation may be written in the form

$$H_{int} = \int dx V(x, t) \rho(x, t) \quad (5)$$

Consider the following expressions for $V(x, t)$, for different time dependence of the potential (spatial form of $V(x)$ remains the same for all cases):

$$V(x, t) = V_0 \theta(t) (1 - \exp(-t/\tau)) \delta(x) \quad (6)$$

$$V(x, t) = V_0 \theta(t) \exp(-t/\tau) \delta(x) \quad (7)$$

$$V(x, t) = V_0 \theta(t) \exp(i\Omega t - \epsilon t) (1 - \exp(-t/\tau)) \delta(x), \quad \epsilon \rightarrow +0 \quad (8)$$

In all cases the potential is nonzero only for positive times due to step function in Eqs.(6-8) and there is no violation of causality.

The matrix element between initial $|i\rangle$ and final $|f\rangle = T_t \exp[i \int_0^t H_{int} dt] |i\rangle$ states in the limit $t \rightarrow \infty$ is given by the following:

$$\langle f|i\rangle = \frac{\int D\varphi \exp[i \int dx dt (S + V(x, t) \rho(x, t))]}{\int D\varphi \exp[i \int dx dt S]}. \quad (9)$$

The integration in Eq.(9) is trivial and we have as a result

$$|\langle f|i\rangle| = \left[\frac{1 + (L/2\pi\tau v)^2}{1 + (1/p_0\tau v)^2} \right]^{-\frac{1}{4\nu} \frac{\delta^2}{\pi^2}}, \quad (10)$$

for the switching on the potential, Eq. (6), and

$$|\langle f|i\rangle| = \left[\frac{1 + (p_0\tau v)^2}{1 + (2\pi\tau v/L)^2} \right]^{-\frac{1}{4\nu} \frac{\delta^2}{\pi^2}}, \quad \delta = -\frac{\nu V_0}{2v} \quad (11)$$

for switching off the potential, Eq.(7), where L is the system length and p_0 is the ultraviolet momentum cut off (of the order of the inverse magnetic length).

In the case of slow switching off ($\tau v \gg L$) the potential or fast switching on ($\tau v \ll L$) we have,

$$|\langle f|i\rangle| \propto L^{-\frac{1}{2\nu} \frac{\delta^2}{\pi^2}}. \quad (12)$$

The time dependence of the matrix element at time $t \gg \tau$ for the potential (6) is easily found:

$$|\langle f(t)|i(0)\rangle| \propto t^{-\frac{1}{\nu} \frac{\delta^2}{\pi^2}}, \quad (13)$$

These results are exact in the infrared limit of the QHE for which the effective Hamiltonian Eq.(4) was derived.

In the case of the periodic potential Eq.(8) $|\langle f|i \rangle| \propto \exp -\Omega/\epsilon \rightarrow 0$, for $\Omega/v < p_0$ and

$$|\langle f|i \rangle| \propto e^{-\frac{1}{2\nu} \frac{\delta^2}{\pi^2} X} \quad (14)$$

for $\Omega > p_0 v$, where $X = 2 \log c + 2(1 - c)/c$ in the limit $\Omega\tau \ll 1$, $X = \frac{(c-1)^2(c+2)}{3c^2(\Omega\tau)^2}$ if $\Omega\tau \gg 1$, and $c = 1 - p_0 v/\Omega$.

The edge states support the nondecaying “particle-hole” pair - the edge wave. This propagating mode with $\chi''(q, \omega) \propto \delta(\omega - vk)$ ($\chi(q, \omega)$ being the density-density correlator) is sufficient to generate the orthogonality in 1D. This would not be the case in higher dimensions, where the existence of the particle-hole continuum, e.g. $\chi''(q, \omega) \propto (\omega/vq)\Theta(\omega - vq)$, with large phase space is required to get orthogonality catastrophe.

Note also that our result Eq.(12) *does not* reproduce the Fermi liquid result at $\nu = 1$. This is the consequence of the chiral nature of the excitations even at $\nu = 1$. To recover the Fermi liquid result, consider the edge of the Fermi liquid state (if one with only edge gapless excitations would exist) as the product of two chiral Slater determinants, each one corresponding to the right and left moving particles. Then Eq.(10) at $\nu = 1$ represents the response of the chiral branch of the Fermi liquid. To get the full answer for the total overlap one has to multiply $\langle f|i \rangle_L \langle f|i \rangle_R = \langle f|i \rangle^2$ and one recovers the standard Fermi liquid result by taking square of the Eq(12) for $\nu = 1$.

Next, we calculate the transition probability for the x-ray problem. Because of the gapped spectrum in the bulk of QH fluid the only gapless excitations to contribute to the x-ray response near frequency threshold are edge excitations. Imagine the x-ray process, creating the deep hole near the edge in a QH sample. Creation of the hole at energy E_0 will lead to the Coulomb potential of the hole and to injection of an electron into the edge of the sample. As is common for the x-ray problem, the response of the edge states will contain two contributions: the orthogonality of the initial and final state Eq(10) and the distortion

of the ground state due to the extra electron. Following the classical paper of Schotte and Schotte [7], we first calculate the propagator:

$$F(t) = \langle i | e^{i \int^t H_i dt} \Psi e^{-i \int^t H_f dt} \Psi^\dagger | i \rangle \quad (15)$$

with H_i being the unperturbed Hamiltonian Eq(1), $H_f = H_i + H_{int}$ is the final state Hamiltonian, and Ψ is the QH liquid electron operator. We assume here that the deep core hole was created in the absorption and is localized. For the case of scatterer in the valence band recoil effects will change our results. Recoil also can be addressed within this bosonization scheme. Using the bosonic algebra we find

$$F(t) \sim t^{-\frac{1}{\nu}(1-\frac{\delta}{\pi})^2}. \quad (16)$$

From this result the transition probability near the frequency edge $W(\omega) \sim \text{Im } i/\pi \int_0^\infty F(t) e^{i(\omega+E_0)t} dt$ becomes:

$$W(\omega) \sim (\omega + E_0)^{\frac{1}{\nu}(1-\frac{\delta}{\pi})^2-1}. \quad (17)$$

In the absence of perturbation V a similar formula was obtained for the electron density of states [8] $N(\omega) \propto \int dk \sum_n | \langle n | \Psi_k | 0 \rangle |^2 \delta(\omega - \omega_n) \propto \omega^{-1+1/\nu}$. The result depends on the potential whose phase shift δ can be either positive or negative. One interesting outcome of this result is that even for fractional QH state the transition probability near the threshold will behave differently for attractive and repulsive interactions. Take for example the unitary phase shift $\delta = \pm\pi/2$ for attractive and repulsive interaction respectively. Then $W(\omega) \sim \omega^{1/(4\nu)-1}$ (*attraction*); $W(\omega) \sim \omega^{9/(4\nu)-1}$ (*repulsion*). And we find even for the $\nu = 1/3$ state that an attractive potential leads to an enhanced transition rate near the threshold: $W(\omega) \sim \omega^{-1/4}$.

To be able to measure the x-ray response of the QH edge, one has to excite a deep hole in the valence-conduction band transition in a semiconductor (with typical bandwidth of the order of 1 eV) and be able to tune the laser to within cyclotron frequency (1-5 meV) to excite only the lowest Landau level and avoid any interlevel excitations. Although the

gapped excitations might be excited in this process, near the threshold frequency, only the soft gapless edge excitations provide the dominant contribution. In the most interesting case of the FQH edge the gaps will be given by the many-body gaps and are somewhat smaller, although of the same magnitude.

There is one interesting aspect of the orthogonality catastrophe in the fractional QH state which makes this problem qualitatively different from the similar problem in a Fermi liquid. In principle one can imagine the “decay” of the electron, injected into the edge, on to fractionally charged quasiparticles, which will provide an extra source of the orthogonalization. However, the simple edge model, used here, does not allow us to address this question. Indeed, in the simple edge model the velocity of the charge on the edge is independent of the charge and therefore the electron will propagate along the edge as a wave packet, which can be considered as made from the quasiparticles. It is only the inclusion of the nonlinear terms and interaction between quasiparticles that might induce the decay of the wave packet.

We can consider the motion of the extra electron taking into account the perturbation potential H_{int} . The matrix element of decay of the electron into $1/\nu$ quasiparticles at points x_i at time t has the form,

$$\begin{aligned} \tilde{F}(t) = & \langle i | \prod_j^{1/\nu} e^{i\varphi(x_j, t)} e^{i \int^t H_{int} dt} e^{-i\varphi(0,0)} | i \rangle \propto \\ & (vt)^{\frac{1}{\nu} \frac{\delta}{\pi} (1 - \frac{\delta}{\pi})} \frac{1}{\prod_j^{1/\nu} (x_j - vt)^{1 - \delta/\pi} x_j^{\delta/\pi}}. \end{aligned} \quad (18)$$

We see that the function $\tilde{F}(t)$ has two maxima : at point $x_j = 0$ and $x_j = vt$. These two peaks have a simple physical origin, the first one near $x = 0$ comes from the screening electron density, induced by the attractive hole potential, while the second one is coming from the propagating mode which moves the electron density along the edge with velocity v . For the case of a strong potential ($\delta = \pi/2$) the probabilities to find the quasiparticles at these points are equal to $(vtp_0)^{-1/(2\nu)}$. Therefore, we assume that the average values of the electric charge at $x = 0$ and at $x = vt$ are the same and equal to $e/2$. Due to the orthogonality catastrophe the matrix element $\tilde{F}(t) \rightarrow 0$ and this effect may be observable

only for $vt < p_0^{-1}$.

Tunneling from metal into QH edge. Next we consider a tunnel junction or point contact between a 1D Fermi liquid and a QH system. Experimentally this situation can be realized by attaching the contact to the edge of the Hall bar.

The Euclidean action of the QH states in terms of a chiral boson field φ follows from Eq. (4) with the substitution $t \rightarrow i\tau$. The noninteracting Fermi liquid can be considered as a superposition of two chiral Fermi liquids (which are described by the action Eq.(4) at $\nu = 1$) with opposite Fermi velocities. The corresponding actions differ by the sign of the $i\partial_\tau\varphi$ term:

$$S_F = S_{\nu=1}(\Psi_L) + S_{\nu=1}(\Psi_R). \quad (19)$$

The tunneling Hamiltonian has the form

$$H_{TUN} = (t_1\Psi_R\Psi^\dagger + t_2\Psi_L\Psi^\dagger)\delta(x) + h.c. \quad (20)$$

Where $\Psi_{L,R}$ are the Fermi liquid left(right) electron operators, Ψ is the electron operator in QH liquid. We do not take into account the spin of the electrons, as we consider only states with spins parallel to the magnetic field B . The opposite spin component can tunnel in the presence of the spin scattering in the contact. Inclusion of this term will simply modify H_{TUN} .

In terms of corresponding boson fields $\varphi_L(x, \tau)$, $\varphi_R(x, \tau)$, $\varphi(x, \tau)$ ($\Psi_{R,L} \propto \exp \pm i\varphi_{R,L}$) we have for the tunneling Hamiltonian

$$\begin{aligned} H_{TUN} = & t_1 \cos(\varphi_R(0, \tau) - \varphi(0, \tau)/\nu) + \\ & t_2 \cos(\varphi_L(0, \tau) + \varphi(0, \tau)/\nu). \end{aligned} \quad (21)$$

Since the perturbation terms act at the one point $x = 0$ it is useful to perform a partial trace in the partition function and integrate in $\varphi, \varphi_R, \varphi_L$ for all $x \neq 0$. If we define $\varphi(0, \tau) = \varphi(\tau)$, $\varphi_{R,L}(0, \tau) = \varphi_{R,L}(\tau)$ then the actions $S_{FQH}, S_{R,L}$ are minimized when

$$\begin{aligned} \varphi(x, \omega_n) = & \varphi(\omega_n), \text{sign}(\omega_n x) \leq 0; \\ = & \varphi(\omega_n) \exp(-\omega_n x), \text{sign}(\omega_n x) > 0 \end{aligned} \quad (22)$$

and similar expressions for φ_R, φ_L . The resulting action is

$$S_{eff} = \frac{1}{4\pi} \left[\frac{1}{\nu} \sum_n |\omega_n| |\varphi(\omega_n)|^2 + \sum_n |\omega_n| |\varphi_R(\omega_n)|^2 + \sum_n |\omega_n| |\varphi_L(\omega_n)|^2 \right] \quad (23)$$

In the lowest order perturbation expansion we obtain the renormalization group (RG) equations for t_1, t_2 upon rescaling $\tau \rightarrow l\tau$

$$\frac{\partial t_{1,2}}{\partial \ln l} = t_{1,2}/2(1 - \nu^{-1}). \quad (24)$$

Both t_1 and t_2 terms flow to zero, indicating that at low energies the junction is insulating. Note that we consider only simple edge states with $\nu \leq 1$. This result can be interpreted as the reflection of the zero overlap between the ground state quasiparticles in the QH liquid with the Fermi liquid quasiparticles, excluding $\nu = 1$ case, where tunneling is not suppressed.

If we cut off the RG flow at some infrared energy scale Λ (corresponding to temperature T , voltage V or frequency), we obtain

$$t_{1,2} \propto \Lambda^{1/2(\nu^{-1}-1)} \quad (25)$$

For $\nu \leq 1$ the conductance of the contact is defined by the largest of $t_{1,2}$ for small Λ . In this case the conductance of the junction is:

$$G \sim t_i^2 \sim V^{\nu^{-1}-1} \quad (26)$$

Therefore we find the power law $I - V$ characteristics, $I \sim V^{\nu^{-1}}$. The nonlinearity in the $I - V$ characteristic is quite strong and, for example, for $\nu = 1/3$ we find $I \sim V^3$, which might be detectable experimentally.

In conclusion, we investigated the influence of time-dependent local perturbations on the ground state of the FQHE state and have calculated the matrix element between the initial and the final ($t \rightarrow \infty$) states. In the limit of sufficiently slow switching off the potential or fast switching on the orthogonality catastrophe occurs. The x-ray problem on the edge of the FQHE state, where the existence of the gapless excitations changes

the response near singularity, is solved. The transition probability for the absorption is $W(\omega) \sim \omega^{\frac{1}{\nu}(1-\frac{\delta}{\pi})^2-1}$. We find that in these problems the phase shift is “fractionalized” $\delta = -\frac{\nu V_0}{2v}$ with filling fraction ν entering into all relevant exponents. We find that for the $\nu = 1/3$ fractional QH state the transition probability is enhanced(suppressed) near the threshold for attractive(repulsive) unitary potential, whereas for any lower filling it will always be suppressed. This is not the case for the lower filling fractions. For the weak potential with $\delta \rightarrow 0$, on the other hand, the transition rate is suppressed for any filling and for both signs of interaction. This situation is *qualitatively* different from the case of Fermi liquid (Eq.(17) with $\nu = 1$), where even in the case of weak potential the sign of the phase shift matters and the transition rate is suppressed(enhanced) for repulsion(attraction). A possible experiment, which probe the x-ray response of the edge might be based on the valence-conduction band transition due to semiconducting laser light absorption, creating a deep hole and electron near the edge. The absorption coefficient will be proportional to the probability $W(\omega)$ Eq. (17). To maximize the absorption at the edges the sample can be covered with only the edges being lightened with laser. If the frequency of the transition is tuned so that gapped bulk excitations are not excited, the dominant response will come from the excited edge states.

We considered the tunnel junction between FQHE system and 1D Fermi liquid as well. We find that for any filling $\nu \leq 1$ the tunneling junction is insulating in the limit $T = 0$ and the nonlinear conductance of the junctions leads to the $I \sim V^{1/\nu}$ tunneling characteristics. We interpret the insulating behavior of the junction as the result of the orthogonality of the electron and the quasiparticles in the QH liquid, except the IQHE case.

Investigation of the dynamical response of the QH states might be an interesting experimental field. We propose to measure the nonlinear tunneling conductance from the QH edge into the metal. If detectable at small bias at low temperature, the $I - V$ characteristics of such a junction will be strongly nonlinear ($I \sim V^3$ for $\nu = 1/3$). The nontrivial dynamics of the edge states can be probed in the experiments with the x-ray transitions in the region near the edge. The x-ray transition will be suppressed or enhanced at the threshold, as

described above.

We are grateful to K. Bedell for comments and careful reading of the manuscript. A.B. would like to thank Prof. H. Jiang for discussion on the possible x-ray transitions experiments in QH liquid. S.M. wishes to thank T-11 and Strongly Correlated Electron Theory program at CMS for hospitality. A.B. thanks M.P.A. Fisher for discussion and for pointing out the mistake in the first version of this paper. This work has been supported in part by the US Department of Energy.

REFERENCES

- [1] B.I. Halperin, Phys. Rev., **B 25**, 2185, (1982).
- [2] X.G. Wen, Int. Journ. of Mod. Phys., **B6**, 1711, (1992).
- [3] C. Kane and M.P.A. Fisher, Phys. Rev., **B 46**, 15233, (1992).
- [4] P.W. Anderson, Phys. Rev. Lett., **18**, 1049, (1967); G.D. Mahan, Phys. Rev., **163**, 612, (1967); P. Nozieres and C.T. De Dominicis, *ibid*, **178**, 1097, (1969).
- [5] See, for example, T. Ogawa, A. Furusaki and N. Nagaosa, Phys. Rev. Lett.**68**, 3638, (1992); D.K.K. Lee and Y. Chen, *ibid*, **69**, 1399, (1992); A. Gogolin, *ibid*, **71**, 2995, (1993) and references therein.
- [6] S. Tomonaga, Prog. Theor. Phys., **5**, 544, (1950).
- [7] K.D. Schotte and U. Schotte, Phys. Rev., **182**, 479, (1969).
- [8] X.G. Wen, Phys. Rev., **B41**, 1238 (1990).